

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Mouse CD48 Antibody[HM48-1]

Catalog Number: E-AB-F1017E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse

Host Armenian Hamster Isotype Armenian Hamster IgG

Clone No. HM48-1

Isotype Control APC Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852E]

Conjugation APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD48 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

Rev. V1.5

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names BCM1 surface antigen;BLAST-1;CD48;CD48 antigen;Cd48;HM48-1;MRC OX-45

surface antigen; SLAMF2; sgp-60

 Uniprot ID
 P18181

 Gene ID
 12506

## For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Background**

CD48 is a 45 kD GPl-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the lg superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

Fax: 1-832-243-6017