

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 647 Anti-Human CD195 Antibody[RoAb13]

Catalog Number: E-AB-F1418M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG2a, κ

Clone No. RoAb13

Isotype Control Elab Fluor[®] 647 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802M]

Conjugation Elab Fluor® 647

Conjugation Information Elab Fluor[®] 647 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with FITC

Anti-Human CD3 Antibody and Elab Fluor[®] 647 Anti-Human CD195 Antibody[RoAb13] (left) or Elab Fluor[®] 647 Mouse IgG2a, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Web: www.elabscience.cn

Shipping Ice bag

Antigen Information

 Uniprot ID
 P51681

 Gene ID
 1234

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD195, also known as CCR5, is a 45 kD G protein-coupled seven transmembrane C C-chemokine receptor. It binds to MIP-1 α , MIP-1 β , and RANTES and is expressed on a subset of T cells and monocytes. CCR5 mediates an intracellular signal thought to induce cell differentiation and proliferation. CCR5 has also been shown to act as a coreceptor for R5 HIV-1 cell entry; modification of CCR5 by sulfation contributes to the efficiency of HIV-1 entry. Studies have shown CCR5 to play a role in a variety of other human diseases, ranging from infectious and inflammatory diseases to cancer.

Web: www.elabscience.cn