A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Mouse IL-17A Antibody[17F3]

catalog number: E-AB-F12720

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Immunogen Recombinant Mouse IL-17A protein

Host Mouse

Isotype Mouse IgGl, κ

Clone 17F3

Purification >98%, Protein A/G purified


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

Applications	Recommended	Dilution

FCM $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-17A gene were stained with 0.2 μg AF/LE Purified Anti-Mouse IL-17A Antibody[17F3] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

Background

The 17F3 monoclonal antibody reacts with mouse IL-17A a 15-20 kDa cytokine expressed by Th17 cells, $\gamma\delta$ T cells, iNKT cells, NK cells, LTi cells, neutrophils, and intestinal Paneth cells. IL-17A has pleiotropic effects in immunoregulation and inflammation. It plays an important role in anti-microbial and chronic inflammation by inducing cytokine and chemokine production, neutrophil influx, and the production of antibacterial peptides but it is also an inflammatory mediator in the development of autoimmune diseases including rheumatoid arthritis, asthma, multiple sclerosis, and psoriasis. The 17F3 antibody has been shown to neutralize IL-18A in vivo.

For Research Use Only

Fax: 1-832-243-6017