

A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse CD30 Protein(His Tag)

Catalog Number: PDMM100076

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Mouse

Source HEK293 Cells-derived Mouse CD30 proteins Phe19-Thr281, C-terminal His

 Calculated MW
 28.8 kDa

 Observed MW
 40-60 kDa

 Accession
 Q60846

Bio-activity Not validated for activity

#### **Properties**

**Purity** > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

#### **Data**



SDS-PAGE analysis of Mouse CD30 proteins, 2  $\mu$ g/lane of Recombinant Mouse CD30 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 28.8 KD

### **Background**

CD30, also known as TNFRSF8, is a cell membrane protein of the tumor necrosis factor receptor family, which regulates proliferation/apoptosis and antibody responses. CD30 is expressed by activated, but not by resting, T and B cells. Aberrant expression of CD30 by mastocytosis mast cells and interaction with its ligand CD30L (CD153) appears to play an important role in the pathogenesis and clinical presentation of systemic mastocytosis. CD30 has been considered as a specific diagnostic biomarker of anaplastic large cell lymphoma (ALCL) and classical Hodgkin lymphoma (cHL). CD30 is also a biomarker used for targeted therapy by an antibody–drug conjugate. Soluble CD30 retains the ability to bind CD30 Ligand and functions as an inhibitor of normal CD30 signaling.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com