

Recombinant Cyclin E2 Monoclonal Antibody

catalog number: E-AB-81550

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	A synthetic peptide of human Cyclin E2
Host	Rabbit
Isotype	Rabbit IgG
Clone	C313336
Purification	Affinity Purified
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.05% stabilizer and 0.05% protective protein.

Applications

Recommended Dilution


WB	1:500-1:1000
IHC	1:50-1:100
IF	1:20-1:100

Data




Western blot detection of Cyclin E2 in Jurkat cell lysates using Cyclin E2 Rabbit mAb(1:500 diluted). Predicted band size: 47kDa. Observed band size: 47kDa.

Observed-MW: 47 kDa
Calculated-MW: 47 kDa



Immunohistochemistry of Cyclin E2 in paraffin-embedded Human lung cancer tissue using Cyclin E2 Rabbit mAb at dilution 1:50



Immunofluorescence of Cyclin E2 (green) in MCF-7 using Cyclin E2 antibody at dilution 1:20, and DAPI(blue)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.9

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2. This cyclin has been shown to specifically interact with CIP/KIP family of CDK inhibitors, and plays a role in cell cycle G1/S transition.