cell system

## 普诺赛® Procell system

## IL-7/LP-1, Mouse, Recombinant

Cat. No.: GPCK352

## 产品信息

**物种** Mouse

表达宿主 E.coli

序列信息 MECHIKDKEGKAYESVLMISIDELDKMTGTDSNCPNNEPNFFRKHVCDDTKEAAFLNRAARKL

cellsystem

KQFLKMNISEEFNVHLLTVSQGTQTLVNCTSKEEKNVKEQKKNDACFLKRLLREIKTCWNKILK

GSI with polyhistidine tag at the C-terminus.

**检索号** P10168.1

标签 His-tag at the C-terminus

**分子量** 15.84 kDa **有效期** 12 months

生物活性 Measured in a cell proliferation assay using PHA-activated human peripheral blood

lymphocytes (PBMC). The ED50 for this effect is < 0.2 ng/mL. The specific activity of

recombinant mouse IL-7 is > 5 × 10<sup>6</sup> IU/mg.

产品特性

**内毒素 (EU/μg)** < 0.1

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored

at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1 × PBS, pH 7.4.

复溶 It is recommended to reconstitute the lyophilized protein in sterile water to a

concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the

biological activity of the protein.

## 背景介绍

Interleukin 7 (IL-7) is a protein that in humans is encoded by the IL7 gene. IL-7 stimulates the differentiation of multipotent (pluripotent) hematopoietic stem cells into lymphoid progenitor cells. It is important for proliferation during certain stages of B-cell maturation, T and NK cell survival, development and homeostasis.

网站: <u>www.procell.com.cn</u> 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址:湖北省武汉市高新大道858号生物医药产业园三期C4栋



