# Recombinant Human XRCC5 & XRCC6 Heterodimer Protein



#### Catalog Number: PKSH030480

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms KARP-1;KARP1;KU80;Ku86;KUB2;NFIV

Species Human

Expression Host Baculovirus-Insect Cells
Sequence Met 1-Ile732&Met 1-Asp609

**Accession** P13010&P12956

Calculated Molecular Weight 85&72 kDa
Observed molecular weight 70 & 85 kDa
Tag N-His

## **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 8.0


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

# **Background**

X-ray repair cross-complementing protein 5, also known as 86 kDa subunit of Ku antigen, ATP-dependent DNA helicase 2 subunit 2, ATP-dependent DNA helicase II 80 kDa subunit, CTC box-binding factor 85 kDa subunit, DNA repair protein XRCC5, Lupus Ku autoantigen protein p86, TLAA and XRCC5, is a nucleus and chromosome which belongs to theku80 family. XRCC5 is a single stranded DNA-dependent ATP-dependent helicase. XRCC5 has a role in chromosome translocation. X-ray repair cross-complementing protein 6, also known as 5'-deoxyribose-5-phosphate lyase Ku70, ATP-dependent DNA helicase 2 subunit 1, ATP-dependent DNA helicase II 70 kDa subunit, 70 kDa subunit of Ku antigen, ATP-dependent DNA helicase 2 subunit 1, CTC box-binding factor 75 kDa subunit, Lupus Ku autoantigen protein p70, Thyroid-lupus autoantigen and XRCC6, is a nucleus and chromosome which belongs to theku70 family. Heterodimer of a

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Human XRCC5 & XRCC6 Heterodimer **Protein**



Catalog Number: PKSH030480

XRCC6 and a XRCC5 subunit associates in a DNA-dependent manner with PRKDC to form the DNA-dependent protein kinase complex DNA-PK, and with the LIG4-XRCC4 complex. The dimer also associates with NAA15, and this complex binds to the osteocalcin promoter and activates osteocalcin expression.

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com