

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Violet 450 Anti-Mouse CD4 Antibody[GK1.5]

Catalog Number: GFH1097UQ

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse Host Rat

Isotype Rat IgG2b, κ
Clone No. GK1.5

Isotype Control Elab Fluor® Violet 450 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843Q]

Conjugation Elab Fluor® Violet 450

**Conjugation Information** Elab Fluor<sup>®</sup> Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).


**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup> cells in 100  $\mu$ L volume].

#### Data



Mouse splenocytes are stained with Elab Fluor<sup>®</sup> Violet 450 Anti-Mouse CD4 Antibody (filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4

Gene ID 12504

### For Research Use Only



## Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

#### **Background**

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, Ick.