# TXNDC12 Polyclonal Antibody

**catalog number:** **E-AB-53059**


**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description


| | |
|---|---|
| **Reactivity** | Human;Mouse;Rat |
| **Immunogen** | Fusion protein of human TXNDC12 |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Purification** | Antigen affinity purification |
| **Buffer** | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|---|---|
| **IHC** | 1:50-1:200 |

## Data



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TXNDC12 Polyclonal Antibody at dilution of 1:60(×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TXNDC12 Polyclonal Antibody at dilution of 1:60(×200)

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| **Shipping** | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

## Background

This gene encodes a member of the thioredoxin superfamily. Members of this family are characterized by a conserved active motif called the thioredoxin fold that catalyzes disulfide bond formation and isomerization. This protein localizes to the endoplasmic reticulum and has a single atypical active motif. The encoded protein is mainly involved in catalyzing native disulfide bond formation and displays activity similar to protein-disulfide isomerases. This protein may play a role in defense against endoplasmic reticulum stress. Alternate splicing results in both coding and non-coding variants.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com