

# Recombinant Human APC protein (His Tag)

Catalog Number: PDEH100955

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Human
<b>Mol_Mass</b>	14.41 kDa
<b>Accession</b>	P25054
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

## Data



SDS-PAGE analysis of Human APC proteins, 2µg/lane of Recombinant Human APC proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 18 KD.

## Background

APC is a ubiquitous, tumor-suppressor protein that is mutated in most colon cancers, generally resulting in a gene product that is truncated in the carboxy terminal region. APC contains several binding domains. The region used as an immunogen overlaps the EBI, DLG (PDZ sequence), and PSD-93 domains. These domains bind microtubules, mitotic spindles, and F-actin and are involved in cell cycle, division, and migration. The immunogen does not include the beta-catenin-binding region, which antagonizes Wnt signaling. The human APC amino acid sequence used as immunogen is 83% and 86% identical to mouse and rat APC, respectively.

## For Research Use Only