

Recombinant RBBP4 Monoclonal Antibody

catalog number: **AN301648L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human RbAp48
Host	Rabbit
Isotype	Rabbit IgG
Clone	C111271
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

WB	1:2000-1:20000
IHC-P	1:1000-1:5000
ICC/IF	1:100-1:200
IP	1:20
FCM	1:20-1:50


Data



Western blot analysis of HeLa, CHO-K1, C6 and Jurkat cell lysates using RBBP4 Antibody (AN301648L) at a dilution of 1:500.

Observed-MW:48 kDa


Calculated-MW:48 kDa




Western blot analysis of Mouse spleen cell lysates using RBBP4 Antibody (AN301648L) at a dilution of 1:1000.

Observed-MW:48 kDa

Calculated-MW:48 kDa



Immunohistochemical analysis of paraffin-embedded Human lung sections using RBBP4 Antibody (AN301648L).



Immunofluorescence analysis of HeLa cells using RBBP4 Antibody(AN301648L). Nuclear DNA was labeled in blue with DAPI. Control: Secondary antibody only.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.5

Background

Core histone-binding subunit that may target chromatin assembly factors, chromatin remodeling factors and histone deacetylases to their histone substrates in a manner that is regulated by nucleosomal DNA. Component of several complexes which regulate chromatin metabolism. These include the chromatin assembly factor 1 (CAF-1) complex, which is required for chromatin assembly following DNA replication and DNA repair; the core histone deacetylase (HDAC) complex, which promotes histone deacetylation and consequent transcriptional repression; the nucleosome remodeling and histone deacetylase complex (the NuRD complex), which promotes transcriptional repression by histone deacetylation and nucleosome remodeling; the PRC2/EED-EZH2 complex, which promotes repression of homeotic genes during development; and the NURF (nucleosome remodeling factor) complex.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.5