Elabscience®

TRPC6 Polyclonal Antibody


catalog number: E-AB-12914


Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | |
|--------------|--|--|
| Reactivity | Human | |
| Immunogen | Synthetic peptide of human TRPC6 | |
| Host | Rabbit | |
| Is otype | IgG | |
| Purification | Affinity purification | |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. | |
| | | |

| Applications | Recommended Dilution |
|--------------|----------------------|
| IHC | 1:50-1:200 |

Data





Immunohistochemistry of paraffin-embedded Human colon Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TRPC6 Polyclonal Antibody at dilution cancer tissue using TRPC6 Polyclonal Antibody at dilution 1:40

1:40

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

The protein encoded by this gene forms a receptor-activated calcium channel in the cell membrane. The channel is activated by diacylglycerol and is thought to be under the control of a phosphatidylinositol second messenger system. Activation of this channel occurs independently of protein kinase C and is not triggered by low levels of intracellular calcium. Defects in this gene are a cause of focal segmental glomerulosclerosis 2 (FSGS2).