
Recombinant Human OMGP/OMG Protein (aa 1-420, His Tag)

Catalog Number: PKSH031717

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human OMGP/OMG protein Met 1-Asn 420, with an C-terminal
	His
Calculated MW	46.0 kDa
Observed MW	120-130 kDa
Accession	P23515-1
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat
	brain glial cells. Immobilized OMG (0.8 μ g/ml, 100 μ l/well) will mediate > 20% C6
	cell adhesion.
Properties	
Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience®

Oligodendrocyte-myelin glycoprotein; also known as OMG and OMGP; is a cell membrane protein which contains eightLRR (leucine-rich) repeats. OMG/ OMGP is a glycosylphosphatidylinositol-anchored protein expressed by neurons and oligodendrocytes in the central nervous system (CNS). OMG/ OMGP is a cell adhesion molecule contributing to the interactive process required for myelination in the central nervous system. OMG/ OMGP play roles in both the developing and adult central nervous system. OMG/ OMGP participats in growth cone collapse and inhibition of neurite outgrowth through its interaction with NgR; the receptor for Nogo. This function requires its leucine-rich repeat domain; a highly conserved region in OMgp during mammal evolution. OMG/ OMGP leucine-rich repeat domain is also implicated in the inhibition of cell proliferation. OMG/ OMGP may also be involved in the formation and maintenance of myelin sheaths. Cell proliferation; neuronal sprouting and myelination are crucial processes involved in brain development and regeneration after injury.