Elabscience®

SNAI1 Polyclonal Antibody

catalog number: E-AB-70119

Note: Centrifuge before opening to ensure complete recovery of vial contents.

ImmunogenKLH conjugated Synthetic peptide corresponding to Mouse SnailHostRabbitIsotypeIgGPurificationAffinity purificationBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol.ApplicationsRecommended DilutionIHC1:500-1:1000		
ImmunogenKLH conjugated Synthetic peptide corresponding to Mouse SnailHostRabbitIsotypeIgGPurificationAffinity purificationBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol.ApplicationsRecommended DilutionHC1:500-1:1000DataImmunohistochemistry analysis of paraffin-embedded mouseImmunohistochemistry analysis of paraffin-embedded mouseImmunohistochemistry analysis of paraffin-embedded mouseImmunohistochemistry analysis of paraffin-embedded Rattestis using SNAI1 Polyclonal Antibody at dilution of 1:600.testis using SNAI1 Polyclonal Antibody at dilution of 1:600.Preparation & StorageStorageStora 4-20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice packupon receipt, store it immediately at the	Description	
Host Rabbit Isotype IgG Purification Affinity purification Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol. Applications Recommended Dilution HC 1:500-1:1000 Data Immunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded Rat testis using SNAI1 Polyclonal Antibody at dilution of 1:600. testis using SNAI1 Polyclonal Antibody at dilution of 1:600. Preparation & Storage Storage Storage Stora at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack.upon receipt,store it immediately at the	Reactivity	Mouse;Rat
sotypeIgGPurificationAffinity purificationBufferPhosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol.ApplicationsRecommended DilutionHC1:500-1:1000DataImmunohistochemistry analysis of paraffin-embedded mousemmunohistochemistry analysis of paraffin-embedded mouseImmunohistochemistry analysis of paraffin-embedded Rat testis using SNAII Polyclonal Antibody at dilution of 1:600. testis using SNAII Polyclonal Antibody at dilution of 1:600.Preparation & StorageStorageStora at -20°C Valid for 12 months. Avoid freeze / thaw cycles.ShippingImmunohistochemistry en immunohistochemistry en immunohistory at the	Immunogen	KLH conjugated Synthetic peptide corresponding to Mouse Snail
Purification Affinity purification Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol. Applications Recommended Dilution IHC 1:500-1:1000 Data Immunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded Rat testis using SNAI1 Polyclonal Antibody at dilution of 1:600. testis using SNAI1 Polyclonal Antibody at dilution of 1:600. Preparation & Storage Storage Storage - 20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping Immunohistochemistry encipation immediately at the	Host	Rabbit
Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol. Applications Recommended Dilution IHC 1:500-1:1000 Data Image: Commended Dilution Image: Commended Dilution Image: Commended Dilution Image: C	Isotype	IgG
Applications Recommended Dilution HC 1:500-1:1000 Data Image: Commended Dilution of 1:600 Image: Commended Dilution of 1:600 Image: Commended Dilution of 1:600 mmunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded Rat testis using SNA11 Polyclonal Antibody at dilution of 1:600 testis using SNA11 Polyclonal Antibody at dilution of 1:600 Preparation & Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack.upon receipt, store it immediately at the	Purification	Affinity purification
Applications Recommended Dilution IHC 1:500-1:1000 Data Image: Constraint of the second s	Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein
IHC 1:500-1:1000 Data Image: Constraint of the second		protectant and 50% glycerol.
Data Image: Data <t< th=""><th>Applications</th><th>Recommended Dilution</th></t<>	Applications	Recommended Dilution
mmunohistochemistry analysis of paraffin-embedded mouse Immunohistochemistry analysis of paraffin-embedded Rat testis using SNAI1 Polyclonal Antibody at dilution of 1:600. testis using SNAI1 Polyclonal Antibody at dilution of 1:600. Preparation & Storage Stor at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	IHC	1:500-1:1000
testis using SNAI1 Polyclonal Antibody at dilution of 1:600. testis using SNAI1 Polyclonal Antibody at dilution of 1:600.Preparation & StorageStorageStorageStore at -20°C Valid for 12 months. Avoid freeze / thaw cycles.ShippingThe product is shipped with ice pack,upon receipt,store it immediately at the	Data	
Preparation & Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack,upon receipt,store it immediately at the	•	
StorageStore at -20°C Valid for 12 months. Avoid freeze / thaw cycles.ShippingThe product is shipped with ice pack,upon receipt,store it immediately at the		
Shipping The product is shipped with ice pack,upon receipt,store it immediately at the		Store at -20°C Valid for 12 months. A void freeze / thaw cycles
	0	-
	Smbhung	

Background

The Drosophila embryonic protein snail is a zinc finger transcriptional repressor which downregulates the expression of ectodermal genes within the mesoderm. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2.