



A Reliable Research Partner in Life Science and Medicine

DUSP13 Polyclonal Antibody

catalog number: E-AB-14067

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Recombinant protein of human DUSP13

Host Rabbit
Isotype IgG


Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

IHC 1:50-1:200

Data





Immunohistochemistry of paraffin-embedded Human thyroid Immunohistochemistry of paraffin-embedded Human colon cancer tissue using DUSP13 Polyclonal Antibody at dilution cancer tissue using DUSP13 Polyclonal Antibody at dilution

1:50

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Members of the protein-tyrosine phosphatase superfamily cooperate with protein kinases to regulate cell proliferation and differentiation. This superfamily is separated into two families based on the substrate that is dephosphorylated. One family, the dual specificity phosphatases (DSPs) acts on both phosphotyrosine and phosphoserine/threonine residues. This gene encodes different but related DSP proteins through the use of non-overlapping open reading frames, alternate splicing, and presumed different transcription promoters. Expression of the distinct proteins from this gene has been found to be tissue specific and the proteins may be involved in postnatal development of specific tissues.