

## Recombinant UCHL3 Monoclonal Antibody

catalog number: **AN300546P**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant Mouse UCHL3 protein
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Clone</b>	4F6
<b>Purification</b>	Protein A
<b>Buffer</b>	0.2 µm filtered solution in PBS


### Applications Recommended Dilution

<b>IHC-P</b>	1:100-1:500
--------------	-------------


### Data



Immunohistochemistry of paraffin-embedded mouse heart using UCHL3 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded mouse brain using UCHL3 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded mouse bone using UCHL3 Monoclonal Antibody at dilution of 1:200.

### Preparation & Storage

<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Shipping</b>	Ice bag

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.0

Ubiquitin carboxyl-terminal hydrolase isozyme L3, also known as UCH-L3, Ubiquitin thioesterase L3 and UCHL3, is a ubiquitin-protein hydrolase that belongs to the peptidase C12 family. It is involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. UCHL3 is highly expressed in heart, skeletal muscle, and testis. UCHL1 and UCHL3 are two of the deubiquitinating enzymes expressed in the brain. These phenotypes indicate the importance of UCHL1 and UCHL3 in the regulation of the central nervous system. UCHL3 functions as a de-ubiquitinating enzyme where lack of its hydrolase activity may result in the prominent accumulation of ubiquitinated proteins and subsequent induction of stress responses in skeletal muscle. UCHL3 has also been identified as a tumor-specific antigen in colon cancer.