A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CD3 epsilon/CD3E (C-Fc-Avi) Biotinylated

Catalog Number: PKSH033930

Note: Centrifuge before opening to ensure complete recovery of vial contents.

	OC	cri	nt	OH
JU		VIII.	34.7	

Species Human

Source HEK293 Cells-derived Human CD3 epsilon; CD3E protein Asp23-Asp126, with an C-

terminal Fc & Avi

Calculated MW 40.5 kDa
Observed MW 45-60 kDa
Accession P07766

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

T-Cell Surface Glycoprotein CD3  $\epsilon$  Chain (CD3 $\epsilon$ ) is a single-pass type I membrane protein. CD3 $\epsilon$  contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3 $\epsilon$  is a polypeptide encoded by the CD3E gene on chromosome 11 in humans. The T cell receptor-CD3 complex (TCR/CD3 complex) is involved in T-cell development and several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs (ITAMs). TCR/CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

### For Research Use Only