

## ECH1 Polyclonal Antibody

catalog number: E-AB-18979

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |  |
|---------------------|--|
| <b>Reactivity</b>   | Human;Rat  |
| <b>Immunogen</b>    | Fusion protein of human ECH1   |
| <b>Host</b>         | Rabbit   |
| <b>Isotype</b>      | IgG  |
| <b>Purification</b> | Antigen affinity purification  |
| <b>Conjugation</b>  | Unconjugated   |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications

### Recommended Dilution

|            |              |
|------------|--------------|
| <b>WB</b>  | 1:500-1:2000 |
| <b>IHC</b> | 1:50-1:300   |

### Data



Western blot analysis of 293T HepG2 and A549 cell lysates using ECH1 Polyclonal Antibody at dilution of 1:200

**Observed-MW:Refer to figures**

**Calculated-MW:36 kDa**



Immunohistochemistry of paraffin-embedded Human tonsil tissue using ECH1 Polyclonal Antibody at dilution of 1:50(×200)

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

This gene encodes a member of the hydratase/isomerase superfamily. The gene product shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species, particularly within a conserved domain characteristic of these proteins. The encoded protein, which contains a C-terminal peroxisomal targeting sequence, localizes to the peroxisome. The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-trans-dienoyl-CoA, indicating that it is a delta3,5-delta2,4-dienoyl-CoA isomerase. This enzyme functions in the auxiliary step of the fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome proliferators.

### For Research Use Only