



A Reliable Research Partner in Life Science and Medicine

PE/Cyanine 5.5 Anti-Mouse CD122/IL-2RB Antibody [5H4]

Catalog Number: E-AB-F1029UI

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ

Clone No. 5H4

PE/Cyanine5.5 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833I] Isotype Control

PE/Cyanine 5.5 Conjugation

Conjugation Information PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

> check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶ cells

in 100 µL volume].

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names IL-2R subunit beta;CD122;High affinity IL-2 receptor subunit beta;IL-2 receptor subunit

beta;IL-2RB;II2rb;Interleukin-2 receptor subunit beta;p70-75

Uniprot ID P16297 Gene ID 16185

Background CD122 is a 70-75 kD IL-2 receptor β chain also known as IL-2Rβ, which is also shared

> by the IL-15 receptor. It is constitutively expressed by NK cells and at lower levels by T cells, B cells, monocytes, and macrophages. The IL-2R\$ chain can combine with either the common y subunit (yc. CD132) alone or with the yc subunit and the IL-2Ra subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation. The 5H4 antibody does not block IL-2 binding to the IL-2 receptor. CD122 is expressed on murine, but not human,

CD8+ Tregs involved in the maintenance of T cell homeostasis.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com