A Reliable Research Partner in Life Science and Medicine

## Recombinant Human CD3D&CD3E Heterodimer (C-Fc-Flag&C-Fc-6His)

Catalog Number: PKSH033951

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$		crip					
	00	0	РΤ	n	ŤΤ		m
v	$\mathbf{c}$	v.		w	w	w	ш

**Species** Human

Source HEK293 Cells-derived Human CD3D&CD3E Heterodimer protein Phe22-Ala105&

Asp23-Asp126, with an C-terminal Fc & Flag & C-terminal Fc & His

 Calculated MW
 37&39.2 kDa

 Observed MW
 40-60 kDa

 Accession
 P04234&P07766

**Bio-activity** Not validated for activity

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

T-cell surface glycoprotein CD3D & CD3E, also known as CD3 delta & CD3 epsilon chain, are single-pass type I membrane proteins. CD3D, together with CD3- epsilon(CD3E), CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.