

Recombinant Rat UCHL3/UCH-L3 Protein (His Tag)

Catalog Number: PKSR030368

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	E.coli-derived Rat UCHL3/UCH-L3 protein Glu 2-Ala 230, with an N-terminal His
Calculated MW	27.5 kDa
Observed MW	28 kDa
Accession	Q91Y78
Bio-activity	Measured by the hydrolysis of UbiquitinAMC. The specific activity is > 14000 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Ubiquitin carboxyl-terminal hydrolase isozyme L3, also known as UCH-L3, Ubiquitin thioesterase L3 and UCHL3, is a ubiquitin-protein hydrolase which belongs to the peptidase C12 family. It is involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. UCHL3 is highly expressed in heart, skeletal muscle, and testis. UCHL1 and UCHL3 are two of the deubiquitinating enzymes expressed in the brain. These phenotypes indicate the importance of UCHL1 and UCHL3 in the regulation of the central nervous system. UCHL3 functions as a de-ubiquitinating enzyme where lack of its hydrolase activity may result in the prominent accumulation of ubiquitinated proteins and subsequent induction of stress responses in skeletal muscle. UCHL3 has also been identified as a tumor-specific antigen in colon cancer.