



## A Reliable Research Partner in Life Science and Medicine

# **IGF2BP3 Polyclonal Antibody**

catalog number: E-AB-62356

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse

**Immunogen** Recombinant fusion protein of human IGF2BP3 (NP 006538.2).


Host Rabbit
Isotype IgG

**Purification** Affinity purification


**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200
IF	1:50-1:200

#### Data




Western blot analysis of extracts of various cell lines using IGF2BP3 Polyclonal Antibody at dilution of 1:1000.




Immunohistochemistry of paraffin-embedded Human esophagus using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).


#### Observed-MV:70 kDa Calculated-MV:21 kDa/63 kDa



Immunohistochemistry of paraffin-embedded Human stomach using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).



Immunohistochemistry of paraffin-embedded Mouse lung using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).



Immunofluorescence analysis of U2OS cells using IGF2BP3 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

#### For Research Use Only

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

**Elabscience®** 

### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

#### Background

The protein encoded by this gene is primarily found in the nucleolus, where it can bind to the 5' UTR of the insulin-like growth factor II leader 3 mRNA and may repress translation of insulin-like growth factor II during late development. The encoded protein contains several KH domains, which are important in RNA binding and are known to be involved in RNA synthesis and metabolism. A pseudogene exists on chromosome 7, and there are putative pseudogenes on other chromosomes.

Fax: 1-832-243-6017