

## IGF2BP3 Polyclonal Antibody

**catalog number: E-AB-62356**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human;Mouse
<b>Immunogen</b>	Recombinant fusion protein of human IGF2BP3 (NP_006538.2).
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications

Applications	Recommended Dilution
<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:50-1:200
<b>IF</b>	1:50-1:200


### Data





Western blot analysis of extracts of various cell lines using IGF2BP3 Polyclonal Antibody at dilution of 1:1000.

**Observed-MW:70 kDa**


**Calculated-MW:21 kDa/63 kDa**



Immunohistochemistry of paraffin-embedded Human stomach using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).



Immunohistochemistry of paraffin-embedded Human esophagus using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).



Immunohistochemistry of paraffin-embedded Mouse lung using IGF2BP3 Polyclonal Antibody at dilution of 1:200 (40x lens).

### For Research Use Only

Immunofluorescence analysis of U2OS cells using IGF2BP3

Polyclonal Antibody at dilution of 1:100. Blue: DAPI for  
nuclear staining.

## Preparation & Storage

**Storage**

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping**

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

The protein encoded by this gene is primarily found in the nucleolus, where it can bind to the 5' UTR of the insulin-like growth factor II leader 3 mRNA and may repress translation of insulin-like growth factor II during late development. The encoded protein contains several KH domains, which are important in RNA binding and are known to be involved in RNA synthesis and metabolism. A pseudogene exists on chromosome 7, and there are putative pseudogenes on other chromosomes.

## For Research Use Only