

Recombinant Rat CLEC4A2/DCIR Protein (His Tag)

Catalog Number: PKSR030276

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	HEK293 Cells-derived Rat CLEC4A2/DCIR protein Gln70-Ser235, with an N-terminal His
Calculated MW	21.8 kDa
Observed MW	32-36 kDa
Accession	Q5YIS1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Dendritic cell immunoreceptor (DCIR), also known as CLEC4A, CLECSF6, is a single-pass type II C-type lectin receptor expressed mainly in dendritic cells (DCs), which is a negative regulator of DC expansion and has a crucial role in maintaining the homeostasis of the immune system. These type II receptors contain a single extracellular carbohydrate recognition domain and have diverse functions in both immunity and homeostasis. DCIR is the only member of the family which contains a cytoplasmic signalling motif and has been shown to act as an inhibitory receptor, while BDCA-2, Dectin-2, DCAR and Mincle all associate with FcRgamma chain to induce cellular activation, including phagocytosis and cytokine production. Dectin-2 and Mincle have been shown to act as pattern recognition receptors for fungi, while DCIR acts as an attachment factor for HIV. In addition to pathogen recognition, DCIR has been shown to be pivotal in preventing autoimmune disease by controlling dendritic cell proliferation. DCIR expressed on antigen presenting cells and granulocytes and acts as an inhibitory receptor via an intracellular immunoreceptor tyrosine-based inhibitory motif (ITIM). It may also be involved via its ITIM motif in the inhibition of B-cell-receptor-mediated calcium mobilization and protein tyrosine phosphorylation. Additionally, DCIR can participate in the capture of HIV-1 and promote infection in trans and in cis of autologous CD4(+) T cells from human immature monocyte-derived DCs. DCIR acts as a ligand for HIV-1 and is involved in events leading to productive virus infection.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017