

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Human CD38 Antibody[AT13/5]

Catalog Number: AN00354E

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. AT13/5

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA


#### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### Data



Staining of normal human peripheral blood cells with APC Anti-Human CD38 Antibody[AT13/5] (filled gray histogram) or APC Mouse IgG1,  $\kappa$  Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Web: www.elabscience.cn

Shipping lce bag

## **Antigen Information**

Alternate Names T10;ADP-ribosyl cyclase

 Uniprot ID
 P28907

 Gene ID
 952

# For Research Use Only

# Elabscience®

# Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

## **Background**

CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.