

A Reliable Research Partner in Life Science and Medicine

APC Anti-Mouse IL-17A Antibody[17F3]

Catalog Number: E-AB-F1272E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. 17F3

APC Mouse IgG1, k Isotype Control[MOPC-21] [Product E-AB-F09792E] Isotype Control

Conjugation

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM**

> of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



HEK293T cells transiently transfected with pcDNA3.1 plasmid encoding Mouse IL-17A gene are stained with APC Anti-Mouse IL-17A Antibody (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Ctla8;II17a;Interleukin-17A

Uniprot ID Q62386 Gene ID 16171

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com Web:www.elabscience.com

Rev. V1.6

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

The 17F3 monoclonal antibody reacts with mouse IL-17A a 15-20 kDa cytokine expressed by Th17 cells, $\gamma\delta$ T cells, iNKT cells, NK cells, LTi cells, neutrophils, and intestinal Paneth cells. IL-17A has pleiotropic effects in immunoregulation and inflammation. It plays an important role in anti-microbial and chronic inflammation by inducing cytokine and chemokine production, neutrophil influx, and the production of antibacterial peptides but it is also an inflammatory mediator in the development of autoimmune diseases including rheumatoid arthritis, asthma, multiple sclerosis, and psoriasis. The 17F3 antibody has been shown to neutralize IL-17A in vivo.

Fax: 1-832-243-6017