A Reliable Research Partner in Life Science and Medicine

Recombinant HER2/ErbB2/CD340 Monoclonal Antibody

catalog number: AN300004P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human


Immunogen Recombinant Human HER2/ErbB2/CD340 Protein

Host Rabbit Isotype lgG Clone A1114 **Purification** Protein A

Buffer 0.2 µm filtered solution in PBS


Applications	Recommended Dilution	
IHC-P	1:100-1:500	
ICC/IF	1:20-1:100	
FCM	1:25-1:100	

Data



Flow cytometric Analysis of HER2/ErbB2/CD340 Monoclonal Immunofluorescence staining of Erbb2 in SKBR3 cells. Cells Antibody on MCF-7 cells. MCF-7 cells were detached using 1× trypsin, washed, then stained with purified rabbit antihuman ErbB2. Second step staining with FITC conjugated goat anti-rabbit IgG (H+L) polyclonal antibody.

were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-Human Erbb2 Monoclonal Antibody (dilution ratio 1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to Cytoplasm and Cell membrane.



Immunohistochemistry of paraffin-embedded human breast carcinoma using HER2/ErbB2/CD340 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

ErbB2, also called Neu and Her2, is a transmembrane glycoprotein in the ErbB family of tyrosine kinase receptors for EGF superfamily growth factors. ErbB2 is widely expressed in epithelial cells and over-expressed in a large number of breast carcinomas. ErbB2 has no identified ligands but heterodimerizes with ErbB1/EGF R, ErbB3, or ErbB4 to form higher affinity signaling complexes. The protease ADAM10 releases a 110 kDa soluble fragment of ErbB2 from the cell surface. ErbB2 plays roles in development, cancer, communication at the neuromuscular junction, and regulation of cell growth and differentiation. The ErbB2/ErbB3 heterodimer is expressed in the majority of breast, skin, ovary and gastrointestinal tumors and transduces a highly mitogenic signal in response to neuregulin 1 (NRG1; heuregulin 1) or NRG2. ErbB3, ErbB2 and neuregulin are all required for formation of the sympathetic nervous system

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2