Elabscience®


PE/Elab Fluor[®] 594 Anti-Human CD19 Antibody[HI19a]

Catalog Number: E-AB-F1304P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgG1, κ
Clone No.	HI19a
Isotype Control	PE/Elab Fluor [®] 594 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792P]
Conjugation	PE/Elab Fluor [®] 594
Conjugation Information	PE/Elab Fluor [®] 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with APC

Anti-Human CD3 Antibody and PE/Elab Fluor[®] 594 Anti-Human CD19 Antibody (Left). Lymphocytes are stained with

APC Anti-Human CD3 Antibody and PE/Elab Fluor[®] 594 Mouse IgG1, κ Isotype Control (Right).

Preparation & Sto	rage
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Informatio	on
Alternate Names	B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>www.elabscience.com</u> Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Uniprot ID	
Gene ID	
Background	

P51677

930

CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cell s, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.