A Reliable Research Partner in Life Science and Medicine

Recombinant Human GM2A Protein (Human Cells, His Tag)

Catalog Number: PKSH033306

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human GM2A protein Ser32-Ile193, with an C-terminal His

Calculated MW 18.6 kDa
Observed MW 19-22 kDa
Accession AAH09273.1

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

 Shipping
 This product is provided as lyophilized powder which is shipped with ice packs.

 Formulation
 Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 7.5.

 Normally 5%
 8% trabaleses mannital and 0.01% Types 80 are added as protestants.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ganglioside GM2 activator (GM2A) is a small glycolipid transport protein which acts as a substrate specific co-factor for the lysosomal enzyme β -hexosaminidase A (HEXB). HEXB together with GM2A, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. GM2A accommodate several single chain phospholipids and fatty acids, is a lipid transfer protein that stimulates the enzymatic processing of gangliosides, and also T-cell activation through lipid presentation. It extracts single GM2 molecules from membranes and presents them in soluble form to beta-hexosaminidase A for cleavage of N-acetyl-D-galactosamine and conversion to GM3.