







AQP2 Polyclonal Antibody


catalog number: E-AB-66725

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human AQP2 (NP_000477.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:50-1:100
IF	1:50-1:200

Data	
 <p>Immunohistochemistry of paraffin-embedded Mouse kidney using AQP2 Polyclonal Antibody at dilution of 1:100 (40x lens).</p>	 <p>Immunofluorescence analysis of Rat kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.</p>
 <p>Immunofluorescence analysis of Mouse kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.</p>	 <p>Immunofluorescence analysis of Rat kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.</p>
 <p>Immunofluorescence analysis of Mouse kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.</p>	 <p>Immunofluorescence analysis of Rat kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.</p>

For Research Use Only



Immunofluorescence analysis of Mouse kidney cells using AQP2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a water channel protein located in the kidney collecting tubule. It belongs to the MIP/aquaporin family, some members of which are clustered together on chromosome 12q13. Mutations in this gene have been linked to autosomal dominant and recessive forms of nephrogenic diabetes insipidus.

For Research Use Only