Recombinant Cynomolgus TPBG/5T4 (C-6His)

Catalog Number: PKSQ050103

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Cynomolgus macaques		
Source	HEK293 Cells-derived Cynomolgus macaques TPBG/5T4 protein Thr35-Ser355, with		
	an C-terminal His		
Calculated MW	35.7 kDa		
Observed MW	55-80 kDa		
Accession	Q4R8Y9		
Bio-activity	Not validated for activity		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	ion Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		
Data			

kDa	МК	R
120 90	_	
60		1002
40		
30	-	
20		
14		

> 95 % as determined by reducing SDS-PAGE.

Background

Trophoblast glycoprotein, also known as TPBG and 5T4, it belongs to the LRR family of proteins. The C-terminus of LRR N-terminal cap (LRRNT) and LRR 1 are essential for the inhibition of the Wnt signaling pathway. TPBG may function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization. It expressed by all types of trophoblasts as early as 9 weeks of development. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma.