
## Recombinant Human FGF basic Protein(Trx Tag)

## Catalog Number: PDEH100657

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description    |  |
|----------------|--|
| Species        | Human  |
| Source         | E.coli-derived Human FGF basic protein Pro143-Ser288, with an N-terminal Trx             |
| Calculated MW  | 36.1 kDa   |
| Observed MW    | 33 kDa   |
| Accession      | P09038   |
| Bio-activity   | Not validated for activity   |
| Properties     |  |
| Purity         | > 95% as determined by reducing SDS-PAGE.  |
| Endotoxin      | < 10 EU/mg of the protein as determined by the LAL method                                |
| Storage        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
|                | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of      |
|                | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.                      |
| Shipping       | This product is provided as lyophilized powder which is shipped with ice packs.          |
| Formulation    | Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%         |
|                | Mannitol.  |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of |
|                | 0.5 mg/mL. Concentration is measured by UV-Vis.  |

## Data



SDS-PAGE analysis of Human FGF basic proteins, 2µg/lane of Recombinant Human FGF basic proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 33 KD

## Background

Basic fibroblast growth factor (bFGF), also known as FGF2, is a member of the fibroblast growth factor (FGF) family. It is a highly specific chemotactic and mitogenic factor for many cell types, appears to be involved in remodeling damaged tissue, such as ulcer healing, vascular repair, traumatic brain injury (TBI). bFGF is a critical component of human embryonic stem cell culture medium. In addition, bFGF protein is a heparin-binding cationic protein involved in a variety of pathological conditions including angiogenesis and solid tumour growth. Thus, bFGF is regarded as a target for cancers chemopreventive and therapeutic strategies.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w v.elabscience.com</u>