Recombinant Human Fas/CD95/TNFRSF6 Protein (His Tag)

Catalog Number: PKSH033436

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Huma	Human	
Source		HEK293 Cells-derived Human Fas/CD95/TNFRSF6 protein Gln26-Asn173, with an C-	
	termir	nal His	
Calculated MW	17.7 kDa		
Observed MW 22-35 kDa		kDa	
Accession	Pession P25445		
Bio-activity	Not v	alidated for activity	
Properties			
Purity > 95 % as		% as determined by reducing SDS-PAGE.	
Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined}$		EU per μ g of the protein as determined by the LAL method.	
Storage Generally, lyophilized prot		ally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Re	econstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	recon	stituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping This product is provided as lyophilized powder which is shippe		product is provided as lyophilized powder which is shipped with ice packs.	
Formulation Lyophilized from a 0.2 μm filtered solution of 20m		nilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.	
	Norm	ally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before	e lyophilization.	
	Please	e refer to the specific buffer information in the printed manual.	
Reconstitution	Please	e refer to the printed manual for detailed information.	
Data			
	kDa MK R		

kDa	MK	R
120 90 60 40 30		-
20	-	
14	-	

> 95 % as determined by reducing SDS-PAGE.

Background

FAS is a receptor and contains three TNFR-Cys repeats and one death domain. It has been shown that FAS is involved in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. FADD (adapter molecule) recruits caspase-8 to the activated receptor, the resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis. FAS-mediated apoptosis may play a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>