

## Recombinant Human TFAP2C/AP2-GAMMA Protein (His Tag)

**Catalog Number:** PKSH030726

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human TFAP2C/AP2-GAMMA protein Leu128-Val223, with an N-terminal His
<b>Calculated MW</b>	12.3 kDa
<b>Observed MW</b>	12-14 kDa
<b>Accession</b>	Q92754
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 98 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 98 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

TFAP2C, also known as AP2-GAMMA, is a member of the activating protein 2 family of transcription factors. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. TFAP2C may be prognostic indicators for patients with breast tumors. TFAP2C gene has been tested for association to diseases (Breast Neoplasms; Carcinoma) and proposed to participate in processes (cell-cell signaling, male gonad development, regulation of transcription from RNA polymerase II promoter). Proteins are expected to have molecular functions (DNA binding, protein binding, protein dimerization activity, transcription factor activity) and to localize in various compartments (membrane, nucleus).