A Reliable Research Partner in Life Science and Medicine

Recombinant Human TFAP2C/AP2-GAMMA Protein (His Tag)

Catalog Number: PKSH030726

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human TFAP2C/AP2-GAMMA protein Leu128-Val223, with an N-

terminal His

 Mol_Mass
 12.3 kDa

 Accession
 Q92754

Bio-activity Not validated for activity

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

TFAP2C, also known as AP2-GAMMA, is a member of the activating protein 2 family of transcription factors. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. TFAP2C may be prognostic indicators for patients with breast tumors. TFAP2C gene has been tested for association to diseases (Breast Neoplasms; Carcinoma) and proposed to participate in processes (cell-cell signaling, male gonad development, regulation of transcription from RNA polymerase II promoter). Proteins are expected to have molecular functions (DNA binding, protein binding, protein dimerization activity, transcription factor activity) and to localize in various compartments (membrane, nucleus).

For Research Use Only

Fax: 1-832-243-6017