



A Reliable Research Partner in Life Science and Medicine

## Recombinant Mouse CXCL1 protein(His Tag)

Catalog Number: PKSM041463

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$					٠.			
H))	es	C	r٦	n	tı	n	m	

**Species** Mouse

Source E.coli-derived Mouse CXCL1 protein Asn 29-Lys 96

 Calculated MW
 7.5 kDa

 Observed MW
 11 kDa

 Accession
 P12850

**Bio-activity** Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The

ED<sub>50</sub> for this effect is <15 ng/mL.

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile 50 mM Tris, 150 mM NaCl, pH 8.5.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Background

Growth-regulated alpha protein (CXCL1,KC), is a member of the alpha chemokine subfamily, was initially identified as an immediate early gene induced in mouse fibroblasts by platelet-derived growth factor. The N-terminal processed form KC(5-72) of the protein is produced by proteolytic cleavage after secretion from bone marrow stromal cells, and shows a highly enhanced hematopoietic activity. Mouse KC shows approximately 63% identity to that of mouse MIP-2. KC is also approximately 60% identical to the human GROs. It has been suggested that mouse KC and MIP-2 are the orthologs of the human GROs and rat CINCs. Cxcl1 has chemotactic activity for neutrophils, and contributes to neutrophil activation during inflammation. Hematoregulatory chemokine, in vitro, suppresses hematopoietic progenitor cell proliferation.