## **RBM15** Polyclonal Antibody

catalog number: E-AB-18870



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description   |  |
|---|--|
| Reactivity  | Human  |
| Immunogen   | Fusion protein of human RBM15  |
| Host  | Rabbit   |
| Isotype   | IgG  |
| Purification  | Antigen affinity purification  |
| Conjugation   | Unconjugated   |
| buffer  | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.                       |
| Applications  | Recommended Dilution   |
| IHC   | 1:50-1:300   |
| Data  |  |
| Immunohistochemistry of paraffin-embedded Human<br>esophagus cancer tissue using RBM15 Polyclonal Antibody<br>at dilution of 1:70(×200)<br>Preparation & Storage  |  |
| - 0   |  |
| Storage   | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping  | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |
| Background  |  |
| RNA-binding motif protein 15 (RBM15) belongs to the SPEN protein family, which have repressor function in several signaling pathway, and may bind to RNA through interaction with spliceosome components. It was first described as a 5' translocation partner of the MAL gene in t(1,22)(p13;q13) infant acute megakaryocytic leukemia, and it functions in murine hematopoiesis through modulating Notch-induced transcriptional activation, which play crucial roles in leukemogenesis. Decreasing RBM15 levels with RNA interference could inhibit the growth and proliferation, block the cell cycle, and induce apoptosis in CML cells. |  |

## For Research Use Only