THG1L Polyclonal Antibody

catalog number: E-AB-53131



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Fusion protein of human THG1L |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:50-1:200 |
| Data | |
| Immunohistochemistry of paraffin-embedded Human tonsil tissue using THG1L Polyclonal Antibody at dilution of 1:50(×200) | |
| Preparation & Storage | |
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |
| Background | |
| The protein encoded by this gene is a mitochondrial protein that is induced by high levels of glucose and is associated with diabetic nephropathy. The encoded protein appears to increase mitochondrial biogenesis, which could lead to renal fibracia. Another function of this protein is that of a guarantee former of diag. CMD to the Stand of tPNA (Uic). Several | |

fibrosis. Another function of this protein is that of a guanyltransferase, adding GMP to the 5' end of tRNA(His). Several transcript variants encoding different isoforms have been found for this gene.

For Research Use Only