## Recombinant Human CXCL11 protein(His Tag)

Catalog Number: PKSH034196



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |   |
|--------------|---|
| Species      | Human   |
| Mol_Mass     | 9.1 kDa   |
| Accession    | O14625  |
| Bio-activity | Measure by its ability to chemoattract BaF3 cells transfected with human CXCR3. The |
|              | ED <sub>50</sub> for this effect is <4 ng/mL.                                       |

|                | $ED_{50}$ for this effect is <4 ng/mL.  |
|----------------|---|
| Properties     |   |
| Purity         | > 98 % as determined by reducing SDS-PAGE.  |
| Endotoxin      | $< 0.1 \; EU$ per µg of the protein as determined by the LAL method.  |
| Storage        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80  |
|                | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| Formulation    | Lyophilized from sterile PBS, pH 7.4.   |
|                | Normally $5\%$ - $8\%$ trehalose, mannitol and $0.01\%$ Tween $80$ are added as protectants   |
|                | before lyophilization.  |
|                | Please refer to the specific buffer information in the printed manual.  |
| Reconstitution | Please refer to the printed manual for detailed information.  |

## Background

CXCL11, also known as I-TAC, SCYB9B, H174 and beta -R1, is a non-ELR CXC chemokine. CXCL11 cDNA encodes a 94 amino acid (aa) residue precursor protein with a 21 aa residue putative signal sequence, which is cleaved to form the mature 73 aa residue protein. CXCL11 shares 36% and 37% amino acid sequence homology with IP-10 and MIG (two other known human non-ELR CXC chemokines), respectively. CXCL11 is expressed at low levels in normal tissues including thymus, spleen and pancreas. The expression of CXCL11 mRNA is radically up regulated in IFN-gamma and I L-1 stimulated astrocytes. Moderate increase in expression is also observed in stimulated monocytes. CXCL11 has potent chemoattractant activity for IL-2 activated T cells and transfected cell lines expressing CXCR3, but not freshly isolated T cells, neutrophils or monocytes. The gene encoding CXCL11 has been mapped to chromosome 4.