# Purified Anti-Human STAT1 Antibody[A15158B]

**catalog number:** **AN010420P**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human |
| **Immunogen** | Recombinant Human STAT1 protein |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Clone** | A15158B |
| **Purification** | >98%, Protein A/G purified |
| **Buffer** | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

| Applications | Recommended Dilution |
|---|---|
| **FCM** | ≤ 0.2 μg per million cells in 100 μL volume |

## Data




A549 cells were fixation and permeabilization, intracellular stained with 0.2 μg Purified Anti-Human STAT1 Antibody[A15158B] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| **Shipping** | Ice bag |

## Background

The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com