A Reliable Research Partner in Life Science and Medicine

Recombinant Human Tie2/CD202b Protein (His Tag)

Catalog Number: PKSH031472

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human Tie2/CD202b protein Met 1-Lys 745, with an C-terminal

His

Calculated MW 82.0 kDa Observed MW 95-105 kDa Accession NP 000450.2

Immobilized recombinant human Tie2 at 2 µg/ml (100 µl/well) can bind human **Bio-activity**

Angiopoietin-2 at a linear range of 1. 28-160 ng/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

This product is provided as lyophilized powder which is shipped with ice packs. Shipping

Lyophilized from sterile PBS, pH 7.4 Formulation


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

TEK, or TIE-2, is an endothelial cell-specific receptor tyrosine kinase (RTK) that is known as a functioning molecule of vascular endothelial cells. TEK comprises a subfamily of RTK with TIE, and these two receptors play critical roles in vascular maturation, maintenance of integrity and remodeling. Targeted mutagenesis of both Tek and its agonistic ligan d, Angiopoietin-1, result in embryonic lethality, demonstrating that the signal transduction pathways mediated by this receptor are crucial for normal embryonic development. TEK signaling is indispensable for the development of the embryonic vasculature and suggests that TEK signaling may also be required for the development of the tumor vasculature.

Fax: 1-832-243-6017