

Purified Anti-Human CD45 Antibody[QA17A19]

catalog number: AN003580P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|---|
| Reactivity | Human |
| Immunogen | Recombinant Human CD45 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | QA17A19 |
| Purification | >98%, Protein A/G purified |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 $\mu\text{g/mL}$ (0.5×10^6 - 1×10^6 cells) |
|------------|---|

Data



Human peripheral blood were stained with 0.2 μg Purified Anti-Human CD45 Antibody[QA17A19] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | Ice bag |

Background

CD45, previously called LCA (leukocyte common antigen), T200, or Ly5 in mice, is member C of the class 1 (receptor-like) protein tyrosine phosphatase family (PTPRC). It is a variably glycosylated 180-220 kDa transmembrane protein that is abundantly expressed on all nucleated cells of hematopoietic origin. Multiple splicing isoforms of exon 4 (A), 5 (B), and 6 (C) are expressed according to cell type, developmental stage and antigenic exposure. The longest form, CD45RABC (called B220 in mouse) is expressed on B lymphocytes, The shortest form, CD45R0, lacking exons 4, 5 and 6 which encode aa 34-194, is expressed on memory cells, while intermediate sizes are expressed on other T cells.

For Research Use Only