PCSK5 Polyclonal Antibody

catalog number: E-AB-52778



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | Description | |
|--|--|--|
| | | |
| Reactivity | Human;Mouse;Rat | |
| Immunogen | Fusion protein of human PCSK5 | |
| Host | Rabbit | |
| Isotype | IgG | |
| Purification | Antigen affinity purification | |
| Conjugation | Unconjugated | |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. | |
| Applications | Recommended Dilution | |
| IHC | 1:25-1:100 | |
| Data | | |
| () shall a shall be | | |
| cancer tissue using PCSk | f paraffin-embedded Human cervical K5 Polyclonal Antibody at dilution of 1:50(×200) | |
| cancer tissue using PCSk Preparation & Storage | X5 Polyclonal Antibody at dilution of 1:50(×200) | |
| cancer tissue using PCSk Preparation & Storage Storage | K5 Polyclonal Antibody at dilution of 1:50(×200) Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. | |
| cancer tissue using PCSk Preparation & Storage | X5 Polyclonal Antibody at dilution of 1:50(×200) | |
| cancer tissue using PCSk Preparation & Storage Storage | K5 Polyclonal Antibody at dilution of 1:50(×200) Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. | |

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER.It then sorts to the trans-Golgi network where a second autocatalytic event takes place and the catalytic activity is acquired. This encoded protein is widely expressed and one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It mediates posttranslational endoproteolytic processing for several integrin alpha subunits and is thought to process prorenin, pro-membrane type-1 matrix metalloproteinase and HIV-1 glycoprotein gp160. Alternative splicing results in multiple transcript variants, some of which encode distinct isoforms, including a protease packaged into dense core granules (PC5A) and a type 1 membrane bound protease (PC5 B).

For Research Use Only