Elabscience®


PE/Cyanine5 Anti-Mouse CD24 Antibody[M1/69]

Catalog Number: E-AB-F1179G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--|---|
| Reactivity | Mouse |
| Host | Rat |
| Isotype | Rat lgG2b, κ |
| Clone No. | M1/69 |
| Isotype Control | PE/Cyanine5 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842G] |
| Conjugation | PE/Cyanine 5 |
| Conjugation Information Storage Buffer | PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter). Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein |
| | protectant. |
| Applications | Recommended usage |
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |





C57BL/6 murine splenocytes are stained with PE/Cyanine5 Anti-Mouse CD24 Antibody[M1/69] (filled gray histogram) or PE/Cyanine5 Rat IgG2b, κ Isotype Control (empty black histogram).

| Preparation & Storage | |
|-----------------------|---|
| Storage | Keep as concentrated solution. |
| Chinaian | This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |
| Antigen Information | |
| Alternate Names | Ly-52;Cd24a;HAS;Nectadrin;R13-Ag;X62 heat stable antigen |
| Uniprot ID | P24807 |
| | |

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Elabscience®

Gene ID Background

12484

CD24 is a 35-45 kD protein also known as Heat Stable Antigen (HSA), Ly-52, or Nectadrin. It is a GPI-linked sialoglycoprotein expressed on lymphocytes, granulocytes, epithelial cells, thymocytes, monocytes, erythrocytes, and dendritic cells. CD24 expression varies during T and B cell differentiation and is a useful marker for delineating various lymphocyte developmental stages. CD24 serves as an adhesion or costimulatory molecule involved in T and B lymphocyte activation and differentiation by homophilic binding or binding to CD62P.

For Research Use Only