# **CD68 Monoclonal Antibody**

catalog number: E-AB-22013



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						
	es	OF	ч г	71	IA	m
ய	$\mathbf{c}$	UI	ш.	յս	ΙU	ш

**Reactivity** Human; Mouse; Rat **Immunogen** Synthetic Peptide

HostMouseIsotypeIgGClone2K6

**Purification** Protein A purification

**Conjugation** Unconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.


Applications Recommended Dilution

**IHC** 1:100-1:300 **IF** 1:50-200


### Data



Immunohistochemistry of paraffin-embedded Human tonsil tissue using CD68 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human tonsil tissue using CD68 Monoclonal Antibody at dilution of 1:200.



Immunofluorescence analysis of Human lung cancer tissue using CD68 Monoclonal Antibody at dilution of 1:200.

## Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

### Background

CD68 is a type I transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It belongs to the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. CD68 is also a member of the scavenger receptor family. It may play a role in phagocytic activities of tissue macrophages.

### For Research Use Only