# Elab Fluor® Red 780 Anti-Mouse CD8a Antibody[53-6.7]

**Catalog Number:** E-AB-F1104US

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*


## Description

| | |
|---|---|
| **Reactivity** | Mouse |
| **Host** | Rat |
| **Isotype** | Rat IgG2a, κ |
| **Clone No.** | 53-6.7 |
| **Isotype Control** | Elab Fluor® Red 780 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833S] |
| **Conjugation** | Elab Fluor®Red 780 |
| **Conjugation Information** | Elab Fluor® Red 780 is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter). |
| **Storage Buffer** | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. |

## Applications — Recommended usage

| | |
|---|---|
| **FCM** | Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/$10^6$ cells in 100 μL volume]. |

## Data




Mouse splenocytes are stained with Elab Fluor® Red 780 Anti-Mouse CD8a Antibody (filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.

## Preparation & Storage

| | |
|---|---|
| **Storage** | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze. |
| **Shipping** | Ice bag |

## Antigen Information

| | |
|---|---|
| **Alternate Names** | CD8A;MAL;T-cell surface glycoprotein CD8 alpha chain;T-lymphocyte differentiation antigen T8/Leu-2 |
| **Uniprot ID** | P01731 |
| **Gene ID** | 12525 |


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com

**Background**

CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form the α(CD8a)/β(CD8b) heterodimer and α/α homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8 α/β heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR α/β T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8 α/α homodimer is expressed on a subset of γ/δ TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase lck.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com