

Recombinant Cynomolgus GITR Ligand/TNFSF18 (C-6His)

Catalog Number: PKSQ050094



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Cynomolgus macaques
Mol_Mass	15.3 kDa
Accession	A0A2K5UCD9
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

TNFSF18 is a single-pass type II membrane protein. It is expressed at high levels in the small intestine, ovary, testis, kidney and endothelial cells. TNFSF18 cytokine binds to TNFRSF18/AITR/GITR. It regulates T-cell responses, and functions as costimulator and lower the threshold for T-cell activation and T-cell proliferation. It is important for interactions between activated T-lymphocytes and endothelial cells and Promotes leukocyte adhesion to endothelial cells. TNFSF18 mediates activation of NF-kappa-B. As Triggers increased phosphorylation of STAT1 and up-regulates expression of VCAM1 and ICAM1. It also regulates migration of monocytes from the splenic reservoir to sites of inflammation.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Tel:400-999-2100

Email:techsupport@elabscience.cn

Web:www.elabscience.cn

Rev. V3.2