

A Reliable Research Partner in Life Science and Medicine

# Recombinant CD79A/Ig-alpha Monoclonal Antibody

catalog number: AN300138P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human; Cynomolgus

Immunogen A synthetic peptide corresponding to the C-terminus of the human CD79a.


HostRabbitIsotypeIgGCloneA1252PurificationProtein A

Buffer 0.2 µm filtered solution in PBS


Applications Recommended Dilution

IHC-P 1:1000-1:20000


#### Data




Immunohistochemistry of paraffin-embedded human tonsil using CD79A / Ig-alpha Monoclonal Antibody at dilution of 1:20000.



Immunohistochemistry of paraffin-embedded human spleen using CD79A / Ig-alpha Monoclonal Antibody at dilution of 1:20000.




Immunohistochemistry of paraffin-embedded human appendix using CD79A / Ig-alpha Monoclonal Antibody at dilution of 1:20000.



Immunohistochemistry of paraffin-embedded cynomolgus spleen using CD79A / Ig-alpha Monoclonal Antibody at dilution of 1:1000.

Rev. V1.3



Immunohistochemistry of paraffin-embedded cynomolgus lymph node using CD79A / Ig-alpha Monoclonal Antibody at dilution of 1:1000.

# **Preparation & Storage**

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com



## **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

**Storage** This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

hinning lee hog

Shipping lce ba

## **Background**

The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (lg). Surface lg non-covalently associates with two other proteins, lg-alpha and lg-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the lg-alpha protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.3