Elabscience®


PE/Elab Fluor[®] 594 Anti-Human CD4 Antibody[RPA-T4]

Catalog Number: E-AB-F1109P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Host	Mouse	
Isotype	Mouse IgG1, κ	
Clone No.	RPA-T4	
Isotype Control	PE/Elab Fluor [®] 594 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792P]	
Conjugation	PE/Elab Fluor [®] 594	
Conjugation Information	PE/Elab Fluor [®] 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).	
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	
Applications	Recommended usage	
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.	

Data



Human peripheral blood lymphocytes are stained with APC

Anti-Human CD19 Antibody and PE/Elab Fluor[®] 594 Anti-Human CD4 Antibody (Left). Lymphocytes are stained with

APC Anti-Human CD19 Antibody and PE/Elab Fluor[®] 594 Mouse IgG1, κ Isotype Control (Right).

Preparation & Stora	ge
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u> Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Uniprot ID	
Gene ID	
Background	

P01730 920

CD4, also known as T4/Leu-3, is a 55 kD single-chain type I transmembrane glycoprotein and member of the immunoglobulin superfamily. It is expressed on most thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part of the TCR/CD3 complex, binds to β 2 domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16 and is a coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (HHV-7).