

IL-23R (C-Fc), Human, Recombinant**Cat. No. : GPCK153****产品信息**

物种	Human
表达宿主	Human Cells
序列信息	Gly24-Asp353
检索号	Q5VWK5
标签	C-Fc
分子量	65.0 kDa
有效期	12 months
生物活性	Immobilized Human IL-23R-Fc at 10 µg/mL (100 µL/well) can bind Human IL-23-His. The ED50 of Human IL-23-His is 0.248 µg/mL.

产品特性

内毒素 (EU/µg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Interleukin 23 Receptor (IL23R) is a type I Cytokine Receptor for IL23. IL23 Receptor complex is comprised of two subunits, the IL12Rβ1 subunit, which is shared with several Cytokines, and a subunit that is unique to IL-23. IL23, after binding to IL23R, activates memory T cells and mediates pro-inflammatory activities in part by the production of IL17 through activation of TH17 lymphocytes. IL23R is expressed on T cells, NK cells, dendritic cells, and macrophages. In fact, polymorphisms of the IL23R gene were reported to be associated with susceptibility to inflammatory diseases and autoimmune diseases such as psoriasis, multiple sclerosis, Graves' s opthalmopathy and inflammatory bowel diseases. The IL23R is known to be critically involved in the carcinogenesis of different malignant tumor.

